



Anti-OGG1 (N-glycosylase/DNA lyase) Acetylated Polyclonal Antibody

Category: Polyclonal Antibody

Catalog #: Acetylated (ac) lysine AB1F059 (Acetylated site: 338K and 341K)

Antigen Synonym: MMH, MUTM, OGH1

Species Reactivity: Human

Immunogen/Specificity:

Polyclonal antibody produced in rabbits immunizing with a synthetic peptide corresponding to C-terminal residues of human OGG1 (N-glycosylase/DNA lyase)

Description: DNA repair enzyme that incises DNA at 8-oxoG residues. OGG1 (N-glycosylase/DNA lyase) excises 7,8-dihydro-8-oxoguanine and 2,6-diamino-4-hydroxy-5-N-methylformamidopyrimidine (FAPY) from damaged DNA. OGG1 has a beta-lyase activity that nicks DNA 3' to the lesion. Defects in OGG1 are associated with tumor formation. Defects in OGG1 are a cause of renal cell carcinoma OGG1 belongs to the type-1 OGG1 family.

Reference:

Aburatani,H., et al, Cancer Res. 57 (11), 2151-2156 (1997)
Rosenquist,T.A., et al, Proc. Natl. Acad. Sci. U.S.A. 94 (14), 7429-7434 (1997)
Roldan-Arjona,T., et al, Proc. Natl. Acad. Sci. U.S.A. 94 (15), 8016-8020 (1997)
Radicella,J.P., et al, Proc. Natl. Acad. Sci. U.S.A. 94 (15), 8010-8015 (1997)
Lu,R., et al, Curr. Biol. 7 (6), 397-407 (1997)
Arai,K., et al, Oncogene 14 (23), 2857-2861 (1997)
Kuo,F.C. and Sklar,J., J. Exp. Med. 186 (9), 1547-1556 (1997)
Bjoras,M., et al, EMBO J. 16 (20), 6314-6322 (1997)

For Research Use Only

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com