



## **Product Information Sheet**

## Biotin Conjugated Goat Anti-mouse IgG

Catalog No.	BA1001-0.5	<i>Purification</i> This antibody was purified from antiserum by immunoaffinity chromatography which removes essentially all goat serum proteins, except the specific antibody for mouse IgG.	
Size	0.5 mg		
lg Type	lgG		
<b>Immunogen</b> Mouse IgG (whole molecular).		Raised in Goat	
Formulation Concentrated, Liquid		<i>Clonality</i> Polyclonal	
Storage Store at $4^{\circ}$ C for frequent use; at $-20^{\circ}$ C for at least one year.		<i>Contents</i> 0.5 mg of biotin conjugated specific antibody. Biotin: Ab=20:1; 0.01M PBS; 1% BSA; 0.01% Thimerosal.	
Avoid multiple freeze-thaw cycles.		Specificity	
Expiration One year from delivery date.		This biotin conjugated antibody is specific for mouse IgG and shows no cross-reactivity with mouse IgA, IgM and other animal species IgG proteins.	
Applications Immunohistochemistry (IHC); Western blot (WB); Direct ELISA		<i>Labeling Method</i> Goat anti-mouse IgG is conjugated to biotin by means of a method described by Bayer. ( <b>Reference</b> : Bayer,E.A. et al.,Meth.Enzymol.,62,308(1979))	
		<i>Application Notes</i> This antibody must be used with various kinds of enzymes and fluorescence	
		conjugated avidin (streptavidin) in IHC, WB and ELISA. <b>Diluted by neutral PBS or TBS.</b>	
То г	reorder contact us at:	Recommended Dilutions	

Antagene, Inc. Toll Free: 1(866)964-2589 Tel: (650) 964-2589 Fax: (650) 964-2519 email: Info@antageneinc.com

## Recommended Dilutions

IHC (paraffin-fixed section)	1:100-200		
IHC (frozen-fixed section)	1:100-200		
WB (DAB coloration)	1:200-400		
Direct ELISA	1:5000-20000		
Optimal working dilutions must be determined by end user.			