



## **Product Information Sheet**

## Biotin Conjugated Rabbit Anti-goat IgG

Catalog No.	BA1006-1	Purification
Size	1 mg	This antibody was purified from antiserum by immunoaffinity chromatography which removes essentially all rabbit serum proteins, except the specific antibody for goat IgG.
Ід Туре	lgG	
Immunogen Goat IgG (whole molecular).		<i>Raised in</i> Rabbit
Formulation Concentrated, Liquid		<i>Clonality</i> Polyclonal
Storage Store at $4^{\circ}$ C for frequent use; at -20 $^{\circ}$ C for at least one year.		<i>Contents</i> 1 mg of biotin conjugated specific antibody. Biotin: Ab=20:1; 0.01M PBS; 1% BSA; 0.01% Thimerosal.
Avoid multiple freeze-thaw cycles.		Specificity
<b>Expiration</b> One year from the day of shipment.		This biotin conjugated antibody is specific for goat IgG and shows no cross-reactivity with goat IgA, IgM and other animal species IgG proteins.
Applications Immunohistochemistry (IHC); Western blot (WB); Direct ELISA		<i>Labeling Method</i> Rabbit anti-goat IgG is conjugated to biotin by means of a method described by Bayer. ( <b>Reference</b> : Bayer,E.A. et al.,Meth.Enzymol.,62,308(1979))
		<i>Application Notes</i> This antibody must be used with various kinds of enzymes and fluorescence conjugated avidin (streptavidin) in IHC, WB and ELISA. <i>Diluted by neutral PBS or TBS.</i>
Toll Free:	reorder contact us at: Antagene, Inc. 1(866)964-2589 Tel: (408)400-9688	Recommended DilutionsIHC (paraffin-fixed section)1:100-200IHC (frozen-fixed section)1:100-200WB (DAB coloration)1:200-400

email: <u>Info@antageneinc.com</u> Website: www.antageneinc.com

Fax: 1(888)225-1868

IHC (paraffin-fixed section)1:100-200IHC (frozen-fixed section)1:100-200WB (DAB coloration)1:200-400Direct ELISA1:5000-20000Optimal working dilutions must be determined by end user.

## FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC AND CLINICAL USE.