



Product Information Sheet

Peroxidase Conjugated Goat Anti-rabbit IgG (γ -chain specific)

Catalog No. BA1054-0.5

Size 0.5 mg

Ig Type IgG

Immunogen

Rabbit IgG (whole molecular).

Storage

Store at 4°C for frequent use; at

-20°C for at least one year.

Avoid multiple freeze-thaw cycles.

Expiration

One year from the day of shipment.

Applications

Dot blot;

Western blot (WB);

Direct ELISA

antiserum by immunoaffinity chromatography which removes essentially all goat serum proteins, except the specific antibody for rabbit IgG.

Raised in Goat

Clonality Polyclonal

Contents

0.5 mg of peroxidase conjugated specific antibody; 0.01M PBS (pH7.4); 1% BSA.

Specificity

This peroxidase conjugated antibody is specific for rabbit IgG and shows no cross-reactivity with rabbit IgA, IgM and other animal species IgG proteins.

Labeling Method

Goat anti-rabbit IgG is conjugated to peroxidase by means of a method described by Wilson MB and Nakane PK.

(**Reference:** Wilson MB and Nakane PK. In Immunofluorescence and Related Staining Techniques, Elsevier/North Holland Biomedical Press, Amsterdam, P215 (1978).)

Preparation of Diluent Buffer

Use 0.01M TBS or 0.01M PBS to dilute. See "Recommended Dilutions" below for details.

Preparation of 0.01M TBS: Add 1.2g Tris, 8.5g NaCl; 450 μ l of purified acetic acid or 700 μ l of concentrated hydrochloric acid to 1000ml H₂O and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Preparation of 0.01 M PBS: Add 8.5g sodium chloride, 1.4g Na₂HPO₄ and 0.2g NaH₂PO₄ to 1000ml distilled water and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Recommended Dilutions

Dot blot (Enhanced chemiluminescent coloration) 1:2000-4000

WB (DAB coloration) 1:500-3000

WB (Enhanced chemiluminescent coloration) 1:3000-10000

Direct ELISA (TMB coloration) 1:20000-40000

Optimal working dilutions must be determined by end user.

To reorder contact us at:

Antagene, Inc.

Toll Free: 1(866)964-2589

Tel: (650) 964-2589

Fax: (650) 964-2519

email: Info@antageneinc.com

Purification

This antibody was purified from