



## Product Information Sheet

### Peroxidase Conjugated Rabbit Anti-goat IgG ( -chain specific)

**Catalog No.** BA1060-0.5

**Size** 0.5 mg

**Ig Type** IgG

**Immunogen**

Goat IgG (whole molecular).

**Storage**

Store at 4°C for frequent use; at

-20°C for at least one year.

Avoid multiple freeze-thaw cycles.

**Expiration**

One year from the day of shipment.

**Applications**

Dot blot;

Western blot (WB);

Direct ELISA

**Purification**

This antibody was purified from antiserum by immunoaffinity chromatography which removes essentially all rabbit serum proteins, except the specific antibody for goat IgG.

**Raised in** Rabbit

**Clonality** Polyclonal

**Contents**

0.5 mg of peroxidase conjugated specific antibody; 0.01M PBS (PH7.4); 1% BSA.

**Specificity**

This peroxidase conjugated antibody is specific for goat IgG and shows no cross-reactivity with goat IgA, IgM and other animal species IgG proteins.

**Labeling Method**

Rabbit anti-goat IgG is conjugated to peroxidase by means of a method described by Wilson MB and Nakane PK.

(**Reference:** Wilson MB and Nakane PK. In Immunofluorescence and Related Staining Techniques, Elsevier/North Holland Biomedical Press, Amsterdam, P215 (1978).)

**Preparation of Diluent Buffer**

Use 0.01M TBS or 0.01M PBS to dilute. See "Recommended Dilutions" below for details.

Preparation of 0.01M TBS: Add 1.2g Tris, 8.5g NaCl; 450µl of purified acetic acid or 700µl of concentrated hydrochloric acid to 1000ml H<sub>2</sub>O and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Preparation of 0.01 M PBS: Add 8.5g sodium chloride, 1.4g Na<sub>2</sub>HPO<sub>4</sub> and 0.2g NaH<sub>2</sub>PO<sub>4</sub> to 1000ml distilled water and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

**Recommended Dilutions**

Dot blot (Enhanced chemiluminescent coloration) 1:2000-4000

WB (DAB coloration) 1:500-3000

WB (Enhanced chemiluminescent coloration) 1:3000-10000

Direct ELISA (TMB coloration) 1:20000-40000

To reorder contact us at:

Antagene, Inc.

Toll Free: 1(866)964-2589

Tel: (650) 964-2589

Fax: (650) 964-2519

email: Info@antageneinc.com

**FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC AND CLINICAL USE.**



# Product Information Sheet

---

*Optimal working dilutions must be determined by end user.*

**FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC AND CLINICAL USE.**

