



Product Information Sheet

Peroxidase Conjugated Protein A

Catalog No. BA1080-1

Size 1 mg

Storage

Store at $4^{\circ}\mathbb{C}$ for frequent use; at $-20^{\circ}\mathbb{C}$ for at least one year. Avoid multiple freeze-thaw cycles.

Expiration

One year from the day of shipment.

Applications

Dot blot;

Western blot (WB);

Direct ELISA

Product Description

Protein A is the component of the cytoderm of staphylococcus aureus. It was regarded as a kind of broad spectrum secondary antibody, because it can bind Fc fragments of a majority of immunoglobulins. It can react with guinea pig, rabbit globulin and human IgG1, 2, 4 strongly and react with mouse, bovine globulin and human IgA, E, M weakly.

Contents

1 mg of peroxidase conjugated protein A; 0.01M PBS (pH7.4); 1% BSA.

Labeling Method

Protein A is conjugated to peroxidase by means of a method described by Wilson MB and Nakane PK.

(**Reference**: Wilson MB and Nakane PK. In Immunofluorescence and Related Staining Techniques, Elsevier/North Holland Biomedical Press, Amsterdam, P215 (1978).)

Preparation of Diluent Buffer

Use 0.01M TBS or 0.01M PBS to dilute. See "Recommended Dilutions" below for details.

Preparation of 0.01M TBS: Add 1.2g Tris, 8.5g NaCl; 450 μ l of purified acetic acid or 700 μ l of concentrated hydrochloric acid to 1000ml H₂O and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Preparation of 0.01 M PBS: Add 8.5g sodium chloride, 1.4g Na_2HPO_4 and 0.2g NaH_2PO_4 to 1000ml distilled water and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

To reorder contact us at:

Antagene, Inc. Recommended Dilutions

Toll Free: 1(866)964-2589 Dot blot (Enhanced chemiluminescent coloration) 1:2000-4000

 Tel: (650) 964-2589
 WB (DAB coloration)
 1:500-3000

 Fax: (650) 964-2519
 WB (Enhanced chemiluminescent coloration)
 1:3000-10000

 email: Info@antageneinc.com
 Direct ELISA (TMB coloration)
 1:20000-40000

Optimal working dilutions must be determined by end user.