



Product Information Sheet

Peroxidase Conjugated Avidin

Catalog No. BA1081-1

Size 1 mg

Storage

Store at 4°C for frequent use; at

-20°C for at least one year.

Avoid multiple freeze-thaw cycles.

Expiration

One year from the day of shipment.

Applications

Dot blot;

Western blot (WB);

Product Description

Avidin is a 68KD protein that extracted from egg white. It has very high affinity to biotin molecule, one million times than the common affinity between antigen and antibody. Avidin is a alkaline protein (IP=10.0-10.5), and it can transfer to be a neutral protein through reconstruction.

Contents

1 mg of peroxidase conjugated avidin; 0.01M PBS (pH7.4); 1% BSA.

Labeling Method

Avidin is conjugated to peroxidase by means of a method described by Wilson MB and Nakane PK.

(**Reference:** Wilson MB and Nakane PK. In Immunofluorescence and Related Staining Techniques, Elsevier/North Holland Biomedical Press, Amsterdam, P215(1978).)

Preparation of Diluent Buffer

Use 0.01M TBS or 0.01M PBS to dilute. See "Recommended Dilutions" below for details.

Preparation of 0.01M TBS: Add 1.2g Tris, 8.5g NaCl; 450μl of purified acetic acid or 700μl of concentrated hydrochloric acid to 1000ml H₂O and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Preparation of 0.01 M PBS: Add 8.5g sodium chloride, 1.4g Na₂HPO₄ and 0.2g NaH₂PO₄ to 1000ml distilled water and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Recommended Dilutions

Dot blot (Enhanced chemiluminescent coloration) 1:2000-4000

WB (DAB coloration) 1:500-3000

WB (Enhanced chemiluminescent coloration) 1:3000-10000

Optimal working dilutions must be determined by end user.

To reorder contact us at:

Antagene, Inc.

Toll Free: 1(866)964-2589

Tel: (650) 964-2589

Fax: (650) 964-2519

email: Info@antageneinc.com

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC AND CLINICAL USE.