



Product Information Sheet

TRITC Conjugated Avidin

Catalog No. BA1093-1

Size 1 mg

Storage

Store at 4 °C. Avoid light and multiple freeze-thaw cycles.

Expiration

One year from the day of shipment.

Applications

Immunocytochemistry (IC); Indirect Immunofluorescence (IF); Immunohistochemistry (IHC).

Product Description

Avidin is a 68KD protein that extracted from egg white. It has very high affinity to biotin molecule, one million times than the common affinity between antigen and antibody. Avidin is a alkalic protein (IP=10.0-10.5), and it can transfer to be a neutral protein through reconstruction.

Contents

1 mg of TRITC conjugated avidin, F/P=3/1; 0.01M PBS; 1% BSA; 0.01% Thimerosal. And the isoelectric point of avidin reduced from 10 to 6.5 through chemical modification. Thus, avidin has very low non-specific absorption to tissue and cell. The emission and filtration wavelength of TRITC are 550 nm and 620 nm respectively.

Labeling Method

Avidin is conjugated to TRITC by means of a method described by Hijmans, W., et al.

(Reference: Hijmans, W., et al. Clin. Exp. Immunol., 4, 457(1969))

Preparation of Diluent Buffer

Add reagent grade BSA into 0.01 M PBS (PH7.2-7.6) or TBS buffer and make BSA at a concentration of 1%. Use the above diluent buffer to dilute. See "Recommended Dilutions" below for details.

Preparation of 0.01M **TBS:** Add 1.2g Tris, 8.5g NaCl; 450 μ l of purified acetic acid or 700 μ l of concentrated hydrochloric acid to 1000ml H₂O and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Preparation of 0.01M **PBS:** Add 8.5g sodium chloride, 1.4g Na₂HPO₄ and 0.2g NaH₂PO₄ to 1000ml distilled water and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

To reorder contact us at:

Antagene, Inc.

Toll Free: 1(866)964-2589

Tel: (650) 964-2589 Fax: (650) 964-2519

email: Info@antageneinc.com

Recommended Dilutions

IC 1: 64
Indirect IF 1: 64
IHC 1: 64

Optimal working dilutions must be determined by end user.