



Product Information Sheet

FITC Conjugated Goat Anti-human IgA

Catalog No. BA1112-0.5

Size 0.5 mg

Immunogen

Human IgA (whole molecular).

Ig Type IgA

Storage

Store at 4 °C. Avoid light and multiple freeze-thaw cycles.

Expiration

One year from the day of shipment.

Applications

Flowcytometry (FCM);

Immunocytochemistry (IC);

Indirect Immunofluorescence (IF);

Immunohistochemistry (IHC).

Purification

This antibody was purified from antiserum by immunoaffinity chromatography which removes essentially all goat serum proteins, except the specific antibody for human IgA.

Raised in Goat

Clonality Polyclonal

Contents

0.5 mg of FITC conjugated specific antibody (purity is above 99%), FITC : Ab = 4-6 : 1; 0.01M PBS; 1% BSA; 0.01% Thimerosal. The emission and filtration wavelength of FITC are 495 nm and 525 nm respectively.

Specificity

This FITC conjugated antibody is specific for human IgA and shows no cross-reactivity with human IgG, IgM and other animal species IgA proteins.

Labeling Method

Goat anti-human IgA is conjugated to FITC by means of a method described by Hijmans, W., et al.

(**Reference:** Hijmans, W., et al. Clin. Exp. Immunol., 4, 457(1969))

Preparation of Diluent Buffer

Add reagent grade BSA into 0.01 M PBS (PH7.2-7.6) or TBS buffer and make BSA at a concentration of 1%. Use the above diluent buffer to dilute. See "Recommended Dilutions" below for details.

Preparation of 0.01M **TBS**: Add 1.2g Tris, 8.5g NaCl; 450µl of purified acetic acid or 700µl of concentrated hydrochloric acid to 1000ml H₂O and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Preparation of 0.01M **PBS**: Add 8.5g sodium chloride, 1.4g Na₂HPO₄ and 0.2g NaH₂PO₄ to 1000ml distilled water and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Recommended Dilutions

FCM 1:100

To reorder contact us at:
Antagene, Inc.
Toll Free: 1(866)964-2589
Tel: (650) 964-2589
Fax: (650) 964-2519
email: Info@antageneinc.com

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC AND CLINICAL USE.

Product Information Sheet

IC	1:16	IHC	1:16
Indirect IF	1:16	<i>Optimal working dilutions must be determined by end user.</i>	