FITC Conjugated Rabbit Anti-human IgG

Catalog No. BA1114-1

Size 1 mg

Ig Type **IgG**

Immunogen

Human IgG (whole molecular).

Storage

Store at 4 °C. Avoid light and multiple freeze-thaw cycles.

Expiration

One year from the day of shipment.

Applications

Flowcytometry (FCM);

Immunocytochemistry (IC);

Indirect Immunofluorescence (IF); Immunohistochemistry (IHC).

Purification

This antibody was purified from antiserum by immunoaffinity chromatography which removes essentially all rabbit serum proteins, except the specific antibody for human IgG.

Raised in Rabbit

Clonality Polyclonal

Contents

1 mg of FITC conjugated specific antibody (purity is above 99%), FITC : Ab = 4-6: 1; 0.01M PBS; 1% BSA; 0.01% Thimerosal. The emission and filtration wavelength of FITC are 495 nm and 525 nm respectively.

Specificity

This FITC conjugated antibody is specific for human IgG and shows no cross-reactivity with human IgA, IgM and other animal species IgG proteins.

Labeling Method

Rabbit anti-human IgG is conjugated to FITC by means of a method described by Hijmans, W., et al.

(Reference: Hijmans, W., et al. Clin. Exp. Immunol., 4, 457(1969))

Preparation of Diluent Buffer

Add reagent grade BSA into 0.01 M PBS (PH7.2-7.6) or TBS buffer and make BSA at a concentration of 1%. Use the above diluent buffer to dilute. See "Recommended Dilutions" below for details.

Preparation of 0.01M TBS: Add 1.2g Tris, 8.5g NaCl; 450µl of purified acetic acid or 700µl of concentrated hydrochloric acid to 1000ml H2O and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Preparation of 0.01M PBS: Add 8.5g sodium chloride, 1.4g Na₂HPO₄ and 0.2g NaH₂PO₄ to 1000ml distilled water and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Recommended Dilutions

To reorder contact us at:

Antagene, Inc. Toll Free: 1(866)964-2589

Tel: (650) 964-2589

Fax: (650) 964-2519

FCM 1:100 IC 1:32 1:32 Indirect IF

Optimal working dilutions must be determined by end user.

1:32

email: Info@antageneinc.com

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC AND CLINICAL USE.

IHC

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC AND CLINICAL USE.