



Product Information Sheet

Polyclonal Anti-Tumor Necrosis Factor α , *TNF α* (Magnetic Bead Conjugate)

Catalogue No. PA1079-M	C-terminal of TNF α of human origin, different from the mouse sequence by one amino acid, and rat sequence by three amino acids.
Lot No. 01010122379125	Purity Immunogen affinity purified.
Ig type: rabbit IgG	Contents Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN ₃ .
Size: 100 μ g/vial	Storage Store at 4°C for frequent use.
Specificity Human, mouse, rat. No cross reactivity with other proteins.	Description This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified magnetic beads. It is useful for immunoprecipitation.
Recommended application Immunoprecipitation(IP)	

Immunogen

A peptide mapping at the

BACKGROUND

Tumor necrosis factor- α (TNF α) also known as TNFR1, It is a potent cytokine, elicits a broad spectrum of biologic responses which are mediated by binding to a cell surface receptor. There are 2 different proteins that serve as major receptors for TNF- α , one associated with myeloid cells and one associated with epithelial cells. TNFR1 maps to 12p13. TNFR1 signaling is also known to activate the transcription factor NF- κ B and promote survival.

REFERENCE

1. Derre, J.; Kemper, O.; Cherif, D.; Nophar, Y.; Berger, R.; Wallach, D. : The gene for the type 1 tumor necrosis factor receptor (TNF-R1) is localized on band 12p13. Hum. Genet. 87: 231-233, 1991.
2. Fuchs, P.; Strehl, S.; Dworzak, M.; Himmler, A.; Ambros, P. F. : Structure of the human TNF receptor 1 (p60) gene (TNFR1) and localization to chromosome 12p13. Genomics 13: 219-224, 1992.
3. Micheau, O.; Tschopp, J. : Induction of TNF receptor I-mediated apoptosis via two sequential signaling complexes. Cell 114: 181-190, 2003.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com