



Product Information Sheet

Polyclonal Anti-MMP14 (Magnetic Bead Conjugate)

Catalogue No. PA1115-M

Immunogen

A synthetic peptide mapping at the C-terminal of human MMP14 different to the related mouse sequence by single amino acid.

Lot No. 08J01

Ig type: rabbit IgG1

Purification

Immunogen affinity purified

Size: 100µg/Vial

Contents

Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN₃.

Specificity

Human, rat, mouse.

No cross reactivity with other proteins.

Storage

Store at 4°C for frequent use.

Recommended application

Immunoprecipitation(IP)

Description:

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified magnetic beads. It is useful for immunoprecipitation

BACKGROUND

The tyrosine phosphatase Shp2 is recruited into tyrosine-kinase signalling pathways through binding of its two amino-terminal SH2 domains to specific phosphotyrosine motifs, concurrent with its re-localization and stimulation of phosphatase activity. Shp2 can potentiate signalling through the MAP-kinase pathway and is required during early mouse development for gastrulation. Shp2 is specifically required in mesenchyme cells of the progress zone (PZ), directly beneath the distal ectoderm of the limb bud. Rather than integrating proliferative signals, Shp2 probably exerts its effects on limb development by influencing cell shape, movement or adhesion. Furthermore, the branchial arches, which also use Fgfs during bud outgrowth, similarly require Shp2. Thus, Shp2 regulates phosphotyrosine-signalling events during the complex ectodermal-mesenchymal interactions that regulate mammalian budding morphogenesis.

REFERENCE

1. Saxton, T. M.; Ciruna, B. G.; Holmyard, D.; Kulkarni, S.; Harpal, K.; Rossant, J.; Pawson, T. : The SH2 tyrosine phosphatase Shp2 is required for mammalian limb development. *Nature Genet.* 24: 420-423, 2000.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com