



Product Information Sheet

Polyclonal Anti-C-C Chemokine receptor 6, **CCR6** (Magnetic Bead Conjugate)

Catalogue No. PA1201-M

Immunogen

Lot No. 08L01

A synthetic peptide corresponding to a sequence at the N-terminal of human CCR6, identical to the related rat and mouse sequence.

Ig type: rabbit IgG1

Purification

Size: 100µg/Vial

Immunogen affinity purified

Specificity

Human, mouse, rat.

No cross reactivity with other proteins.

Contents

Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN₃.

Storage

Store at 4°C for frequent use.

Recommended application

Immunoprecipitation(IP)

Description:

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified magnetic

BACKGROUND

CCR6, also termed CKRL3, encodes a 369-amino acid polypeptide with greatest similarity to the family of alpha-chemokine-binding receptors¹. Unlike most chemokine receptor genes, it is encoded by more than 1 exon². Weakly expressed as a 4-kb transcript in spleen, lymph nodes, peripheral blood lymphocytes and appendix, CCR6 gene is located at 6q27. As the receptor for MIP-3-alpha, its activation leads to phospholipase C-dependent intracellular Ca(2+) mobilization. Additionally, CCR6 are markedly upregulated in psoriasis³.

REFERENCE

1. Zaballo, A.; Varona, R.; Gutierrez, J.; Lind, P.; Marquez, G. : Molecular cloning and RNA expression of two new human chemokine receptor-like genes. *Biochem. Biophys. Res. Commun.* 227: 846-853, 1996.
2. Liao, F.; Lee, H.-H.; Farber, J. M. : Cloning of STRL22, a new human gene encoding a G-Protein-coupled receptor related to chemokine receptors and located on chromosome 6q27. *Genomics* 40: 175-180, 1997.
3. Homey, B.; Dieu-Nosjean, M.-C.; Wiesenborn, A.; Massacrier, C.; Pin, J.-J.; Oldham, E.; Catron, D.; Buchanan, M. E.; Muller, A.; de Waal Malefyt, R.; Deng, G.; Orozco, R.; Ruzicka, T.; Lehmann, P.; Lebecque, S.; Caux, C.; Zlotnik, A. : Up-regulation of macrophage inflammatory protein-3-alpha/CCL20 and CC chemokine receptor 6 in psoriasis. *J. Immun.* 164: 6621-6632, 2000.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com