



Product Information Sheet

Polyclonal Anti- matrix metalloproteinase-10, MMP-10 (Magnetic Bead Conjugate)

Catalogue No. PA1380-M Lot No. 0131112098027	Immunogen A synthetic peptide corresponding to a sequence at the C-terminal of human MMP-10 (410-422 aa), identical to the related mouse and rat sequence.
Ig type rabbit IgG	Purity Immunogen affinity purified.
Size 100µg/vial	
Specificity Human. No cross reactivity with other	Contents Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na $_2$ HPO $_4$, 0.05mg Thimerosal, 0.05mg NaN $_3$.
proteins. Recommended application	Reconstitution 0.2ml of distilled water will yield a concentration of 500µg/ml.
ImmunoPrecipitation	Storage At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for longer time.

BACKGROUND

Stromelysin-2 also known as matrix metalloproteinase-10 (MMP-10) or transin-2 is an enzyme that in humans is encoded by the MMP10 gene.Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades proteoglycans and fibronectin. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3.

REFERENCE

- 1.Muller D, Quantin B, Gesnel MC, Millon-Collard R, Abecassis J, Breathnach R (July 1988). "The collagenase gene family in humans consists of at least four members". Biochem. J. 253 (1): 187–92.
- 2.Jung JY, Warter S, Rumpler Y (1990). "Localization of stromelysin 2 gene to the q22.3-23 region of chromosome 11 by in situ hybridization". Ann. Genet. 33 (1): 21–3.