



## **Anti- TDG (G/T mismatch-specific thymine DNA glycosylase) Ubiquitinated Polyclonal Antibody**

**Category:** Polyclonal Antibody

**Catalog #:** monoubiquitinated lysine **AB1F065 (Site: K330)**

**Species Reactivity:** Human, Mouse, rat

### **Immunogen/Specificity:**

Polyclonal antibody produced in rabbits immunizing with a synthetic peptide corresponding to C-terminal residues of human TDG (G/T mismatch-specific thymine DNA glycosylase)

Note: \* At K330 amino acid residue with the ubiquitin c-terminal 7-mer peptide bound: CLRLRGG-

**Description:** In the DNA of higher eukaryotes, hydrolytic deamination of 5-methylcytosine to thymine leads to the formation of G/T mismatches. TDG(G/T mismatch-specific thymine DNA glycosylase) corrects G/T mispairs to G/C pairs. It is capable of hydrolyzing the carbon-nitrogen bond between the sugar-phosphate backbone of the DNA and a mispaired thymine. In addition to the G/T, TDG can remove thymine also from C/T and T/T mispairs in the order G/T >> C/T > T/T. It has no detectable activity on apyrimidinic sites and does not catalyze the removal of thymine from A/T pairs or from single-stranded DNA. TDG can also remove uracil and 5-bromouracil from mispairs with guanine.

Sumoylation on Lys-330 by either SUMO1 or SUMO2 induces dissociation of the product DNA.

TDG belongs to the TDG/mug DNA glycosylase family.

### **Reference:**

Neddermann,P., et al, J. Biol. Chem. 271 (22), 12767-12774 (1996)

Neddermann,P. and Jiricny,J., Proc. Natl. Acad. Sci. U.S.A. 91 (5), 1642-1646 (1994)

Hardeland,U., et al, EMBO J. 21 (6), 1456-1464 (2002)

Baba,D., et al, Nature 435 (7044), 979-982 (2005)

**For Research Use Only**

**Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: [Info@antageneinc.com](mailto:Info@antageneinc.com)**